


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



NPC-CMAPSEEC Issue

Volume 12, Issue 2, Pages 153-318, 2017
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS**PROFESSOR ALEJANDRO F. BARRERO**

Department of Organic Chemistry, University of Granada,
Campus de Fuente Nueva, s/n, 18071, Granada, Spain
afbarre@ugr.es

PROFESSOR MAURIZIO BRUNO

Department STEBICEF,
University of Palermo, Viale delle Scienze,
Parco d'Orleans II - 90128 Palermo, Italy
maurizio.bruno@unipa.it

PROFESSOR VLADIMIR I. KALININ

G.B. Elyakov Pacific Institute of Bioorganic Chemistry,
Far Eastern Branch, Russian Academy of Sciences,
Pr. 100-letiya Vladivostoka 159, 690022,
Vladivostok, Russian Federation
kalininv@piboc.dvo.ru

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimakiy@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry, University of Wollongong,
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry, Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry, The University of Alabama in Huntsville,
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR PING-JYUN SUNG

National Museum of Marine Biology and Aquarium
Checheng, Pingtung 944
Taiwan
pjsung@nmmba.gov.tw

PROFESSOR YASUHIRO TEZUKA

Faculty of Pharmaceutical Sciences, Hokuriku University,
Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan
y-tezuka@hokuriku-u.ac.jp

PROFESSOR DAVID E. THURSTON

Institute of Pharmaceutical Science
Faculty of Life Sciences & Medicine
King's College London, Britannia House
7 Trinity Street, London SE1 1DB, UK
david.thurston@kcl.ac.uk

HONORARY EDITOR**PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Giovanni Appendino
Novara, Italy

Prof. Norbert Arnold
Halle, Germany

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Vassaya Bankova
Sofia, Bulgaria

Prof. Roberto G. S. Berlinck
São Carlos, Brazil

Prof. Anna R. Bilia
Florence, Italy

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Fatih Demirci
Eskişehir, Turkey

Prof. Francesco Epifano
Chieti Scalo, Italy

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Dr. Christopher Gray
Saint John, NB, Canada

Prof. Dominique Guillaume
Reims, France

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Hisahiro Hagiwara
Niigata, Japan

Prof. Judith Hohmann
Szeged, Hungary

Prof. Tsukasa Iwashina
Tsukuba, Japan

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Phan Van Kiem
Hanoi, Vietnam

Prof. Niel A. Koorbanally
Durban, South Africa

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Lacaillle-Dubois
Dijon, France

Prof. Shoei-Sheng Lee
Taipei, Taiwan

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Stefano Serra
Milano, Italy

Dr. Bikram Singh
Palampur, India

Prof. Leandros A. Skaltsounis
Zografou, Greece

Prof. John L. Sorensen
Manitoba, Canada

Prof. Johannes van Staden
Scottsville, South Africa

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2017 subscription price: US\$2,595 (Print, ISSN# 1934-578X); US\$2,595 (Web edition, ISSN# 1555-9475); US\$2,995 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Phytochemical Profile of *Inula britannica* from Bulgaria

Victoria Ivanova^a, Antoaneta Trendafilova^{a,*}, Milka Todorova^{a,*}, Kalina Danova^a and Dimitar Dimitrov^b

^aInstitute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Acad. G. Bonchev str., bl. 9, 1113 Sofia, Bulgaria

^bNational Museum of Natural History, Bulgarian Academy of Sciences, 1, Tsar Osvoboditel Blvd., 1000 Sofia, Bulgaria

trendaf@orgchm.bas.bg; todorova@orgchm.bas.bg

Received: June 2nd, 2016; Accepted: July 29th, 2016

The flower heads of *Inula britannica* L. of Bulgarian origin afforded sesquiterpene lactones (gaillardin, britannin, 11,13-dihydroinuchinenolide B, ivalin, pulchellin C), triterpenoids (3-*O*-palmitates of 16 β -hydroxylupeol, 16 β -hydroxy- β -amyrin, and faradiol) and flavonoids (quercetin, luteolin, luteolin-7-*O*-glucoside). All compounds are known and they were identified by spectral methods. The observed differences in the chemical content of the chloroform and methanol extracts were also reflected in their free radical scavenging activity, evaluated by DPPH and ABTS assays. Intraspecific variability of *I. britannica* is discussed.

Keywords: *Inula britannica*, Asteraceae, Sesquiterpene lactones, Triterpenoids, Flavonoids, DPPH and ABTS assays.

The genus *Inula* (Asteraceae) consists of approximately 100 species, distributed mainly in Asia and Europe. Many studies have been carried within the genus due to the significant structural diversity and biological activity of sesquiterpene lactones, diterpenes, triterpenes, and flavonoids, isolated from a number of *Inula* species [1,2]. *I. britannica* is used in Chinese folk medicine to treat digestive disorders, bronchitis and inflammation, bacterial and viral infections, as well as some tumors [2b]. That is why most of the studies are on *I. britannica* of Chinese origin. The application in traditional medicine requires extensive phytochemical study and pharmacological evaluations of this species. It should be noted that sesquiterpene lactones were the largest group. More than 40 lactones of different skeletal types – germacranolides, eudesmanolides, guaianolides, secoeudesmanolides, seco-guaianolides, and lactone dimers, have been described so far. To the best of our knowledge, from the European taxa only Russian [3] and Serbian [4] have been investigated up to now.

In continuation of our studies on species of the Asteraceae family this work is a part of a phytochemical investigation of Bulgarian taxa from the genus *Inula*. The CHCl₃ extract of the flowers of *I. britannica* was fractionated by column chromatography on silica gel. IR control allowed selection of the fractions (characteristic absorption band at 1770-1740 cm⁻¹) for further determination of the lactone profile. Thus, the sesquiterpene lactones ivalin [5], britannin [3a,4], gaillardin [4,5a], 11,13-dihydroinuchinenolide B [4], and pulchellin C [3b,4] have been isolated and identified. All these compounds have been previously found in *I. britannica* of different origins. With exception of ivalin, the other compounds were reported for the Serbian taxon [4]. From the latter one, 1,10-secoeudesmanolides – 14-(3-methylpentanoyl)-6-deoxybritannilactone, 14-(3-methylbutanoyl)-6-deoxybritannilactone, and 14-(2-methylpropanoyl)-6-deoxybritannin, as well as 4H-tomentosin have been isolated. Varieties of lactones with secoeudesmane and secoguaiane skeletons have been reported before in the Asian populations of *I. britannica* [1,2]. It should be noted that no one of these substances was detected in the studied Bulgarian sample.

It was found (IR control) that the less polar fractions did not contain sesquiterpene lactones, but the absorption band at 1720 cm⁻¹ showed the presence of compounds bearing ester groups. Further on, prep. TLC yielded the known triterpenoids 3-*O*-palmitates of 16 β -hydroxylupeol, 16 β -hydroxy- β -amyrin, and faradiol. Only the first one has been found in a Turkish population of *I. britannica* so far [6a], while the other two compounds are isolated now for the first time from this species, but they have been detected earlier in *Achillea alexandri-regis* [6b] and *Calendula officinalis* [6c], respectively. β -Sitosterol and β -amyrin were also detected. More or less, different triterpenoids have been isolated from the genus *Inula*, but esters of long-chain fatty acids have been detected only in one Turkish population of *I. britannica* [6a]. It should be noted that the literature data for triterpenoids in *Inula* species are insufficient for discussion of their chemotaxonomic significance.

Besides the described above compounds, 3 flavonoids and 1,5-dicafeloylquinic acid were isolated from the MeOH extract. Luteolin, luteolin-7-*O*-glucoside and quercetin are known for *I. britannica*, while 1,5-dicafeloylquinic acid was detected in *I. viscosa* [2]. Being very common plant components and usually in high concentrations, at this point of the investigation, these compounds were not regarded as compounds of chemotaxonomic interest.

Further, TLC comparison of extracts from flowers and leaves showed that both plant organs accumulated the same sesquiterpene lactones. Regarding phenolic components, it was found that luteolin, luteolin-7-*O*-glucoside, and 1,5-dicafeloylquinic acid were present in both leaves and flowers, while quercetin was detected only in flowers. Finally, the observed differences in the chemical composition of the chloroform and methanol extracts were also reflected in their free radical scavenging activity, evaluated by DPPH and ABTS assays (Table 1). As can be seen, both methanol extracts (from flowers and leaves) possessed the highest antioxidant activity and maximal total phenolic and flavonoid contents. A good correlation between antiradical activity in the studied extracts and their total phenolic content was also observed ($R^2 = 0.9954$ and $R^2 = 0.9910$ for ABTS and DPPH, respectively).

Table 1: Total phenolic (TPC) and flavonoid (FC) contents in different *I. britannica* extracts and their radical scavenging activity.

Plant parts	Extract	TPC	FC	ABTS	DPPH
		[mgGA/gDM]	[mgC/gDM]	[μ MT/gDM]	[μ MT/gDM]
Leaves	CHCl ₃	0.4±0.02	0.3±0.02	1.6±0.1	0.7±0.1 ^a
Flowers	CHCl ₃	0.1±0.02	0.1±0.02	1.1±0.1	0.7±0.02 ^a
Leaves	MeOH	2.3±0.1	0.6±0.01	15.5±0.2	13.7±0.7
Flowers	MeOH	7.9±0.4	0.9±0.02	44.4±0.4	37.6±0.6

^aValues with the same letter are not significantly different, $p \leq 0.05$. Standard deviations were calculated on the base of three samples.

Diversity of lactones found in the so far studied taxa of *I. britannica* of Asian and European origin revealed significant intraspecific variability. Nevertheless, on the basis of the skeleton type of lactones it could be suggested that secoeudesmanolides and secoguaianolides characterized the Asian populations. On the other hand, Bulgarian and Russian (European) populations are free of these types of lactones. Further, secoeudesmanolides and one secoguaianolide have been isolated from a Serbian taxon, but bicyclic lactones are principal components. So, the existence of secoeudesmanolides and secoguaianolides in the European *I. britannica* cannot be ruled out. Thus, the domination of guaianolides and eudesmanolides or their seco-derivatives characterize European and Asian taxa, respectively. Further phytochemical investigation on *I. britannica* will clarify the lactone profile of European populations.

Experimental

Plant material: *I. britannica* was collected from a natural locality in the Southern Balkan Region in Bulgaria. A voucher specimen (SOM 172474) was deposited in the Herbarium of the Institute of Biodiversity and Ecosystem Research, Bulg. Acad. Sci.

Extraction and isolation: A portion (1 g) of air-dried flower heads and leaves of *I. britannica* were extracted with CHCl₃ and MeOH. The corresponding crude extracts were compared by TLC (Silica gel 60 and RP-18, F₂₅₄ (Merck), CHCl₃-diethyl ether, 50:1; CHCl₃-acetone, 10:1 and 1:1; MeOH-H₂O, 1:1) and used for determination of the total phenolics, flavonoids and antioxidant capacity. Further, the CHCl₃ and MeOH extracts from flower heads (30 g) of the plant were worked up for isolation of the individual compounds. The CHCl₃ extract (1.5 g) was fractionated by column chromatography (CC) on silica gel using CHCl₃-acetone mixtures with increasing

polarity to give 11 fractions (F₁-F₁₁). Further separation by CC and prep. TLC (silica gel, *n*-hexane-diethyl ether, 5:1) of Fr. F₂ (50 mg) afforded 3-*O*-palmitates of 16 β -hydroxyulpeol (5.7 mg), 16 β -hydroxy- β -amyryn (5.3 mg), and faradiol (5.0 mg). The presence of β -amyryn and β -sitosterol was proved by TLC of F₃ (silica gel, *n*-hexane-diethyl ether, 1:1) using these compounds as standards. Prep. TLC (silica gel, CHCl₃-acetone, 5:1) of F₇ (47 mg) afforded ivalin (7 mg) and britannin (14 mg). Gaillardin (33 mg) and pulchellin C (18 mg) were obtained from F₉ (65 mg) and F₁₁ (92 mg), resp. after recrystallization (CHCl₃). Prep. TLC (silica gel, CHCl₃-acetone, 5:1) of F₁₀ (18 mg) yielded 11,13-dihydroinuchinenolide B (7 mg). The MeOH extract (0.5 g) was separated into 2 fractions F-1 and F-2 by CC (Sephadex LH-20, MeOH). Prep. TLC (silica gel RP-18, MeOH-H₂O, 1:1) of F-2 (50 mg) yielded luteolin (9 mg), quercetin (1.8 mg), luteolin-7-*O*-glucoside (2.0 mg), and 1,5-dicafeyloylquinic acid (6.4 mg). All isolated compounds were proved by comparison of their ¹H NMR spectral data with those in the literature.

Determination of the content of total phenolic compounds (TPC) and total flavonoids (FC): TPC and FC were determined by procedures described in ref. [7a,b] and expressed as mg gallic acid equivalents per g dry plant material [mgGA/gDM] and mg (+)-catechin equivalents per g dry plant material [mgC/gDM], respectively.

DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging activity and ABTS [2,2'-azino-bis(3-ethylbenzthiazoline-6-sulphonic acid)] assays: The DPPH and ABTS assays were performed according to the procedures described by Thaipong *et al.* [7c] and the antioxidant activity was expressed as μ M Trolox equivalents per g dry plant material [μ MT/gDM].

Statistical analysis: Correlation coefficients (R^2) for determination of the relationship between the radical scavenging activity and the TPC and FC were calculated using MS Excel software.

Acknowledgments - This work was supported by the framework of the Bulgarian-Swiss Research Programme under Grant No. IZEBZ0_142989; DO2-1153).

References

- [1] (a) Khan AL, Hussain J, Hamayun M, Gilani SA, Ahmad S, Rehman G, Kim YH, Kang SM, Lee IJ. (2010) Secondary metabolites from *Inula britannica* L. and their biological activities. *Molecules*, **15**, 1562-1577; (b) Wang GW, Qin JJ, Cheng XR, Shen YH, Shan L, Jin HZ, Zhang WD. (2014) *Inula* sesquiterpenoids: structural diversity, cytotoxicity and anti-tumor activity. *Opinion on Investigational Drugs*, **23**, 318-345.
- [2] (a) Seca AML, Grigore A, Pinto DCGA, Silva AMS. (2014) The genus *Inula* and their metabolites: From ethnopharmacological to medicinal uses. *Journal of Ethnopharmacology*, **154**, 286-310; (b) Seca AML, Pinto DCGA, Silva AMS. (2015) Metabolomic profile of the genus *Inula*. *Chemistry & Biodiversity*, **12**, 859-906.
- [3] (a) Rybalko K, Sheichenko V, Maslova G, Kiseleva E, Gubanov I. (1968) Britannin – a lactone from *Inula britannica*. *Chemistry of Natural Compounds*, **4**, 215-215; (b) Serkerov S, Mir-Babaev N. (1988) Pulchellin C from *Inula britannica*. *Chemistry of Natural Compounds*, **24**, 752-753.
- [4] Fishedick JT, Pesic M, Podolski-Renic A, Bankovic J, de Vos RCH, Peric M, Todorovic S, Tanic N. (2013) Cytotoxic activity of sesquiterpene lactones from *Inula britannica* on human cancer cell lines. *Phytochemistry Letters*, **6**, 246-252.
- [5] (a) Ito K, Iida T. (1981) Seven sesquiterpene lactones from *Inula britannica* var. *chinensis*. *Phytochemistry*, **20**, 271-273; (b) Yang JL, Wang R, Liu LL, Shi YP. (2011) Sesquiterpenoids from *Inula britannica*. *Planta Medica*, **77**, 362-367.
- [6] (a) Oksuz S, Topcu G. (1987) Triterpene fatty acid esters and flavonoids from *Inula britannica*. *Phytochemistry*, **26**, 3082-3084; (b) Kundakovic T, Fokialakis N, Magiatis P, Kovacevic N, Chinou I. (2004) Triterpene derivatives of *Achillea alexandri-regis* BORN. & RUDSKI. *Chemical & Pharmaceutical Bulletin*, **52**, 1462-1465; (c) Eitterl-Eglseer K, Reznicek G, Jurenitsch J, Novak J, Zitterl W, Franz C. (2001) Morphogenetic variability of faradiol monoesters in marigold *Calendula officinalis* L. *Phytochemical Analysis*, **12**, 199-201.
- [7] (a) Yoo KM, Lee CH, Lee H, Moon BK, Lee CY (2008) Relative antioxidant and cytoprotective activities of common herbs. *Food Chemistry*, **106**, 929-936; (b) Zhishen J, Mengcheng T, Jianming W. (1999) The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry*, **64**, 555-559; (c) Thaipong K, Boonprakob U, Crosby K, Cisneros-Zevallos L, Byrne DH. (2006) Comparison of ABTS, DPPH, FRAP and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis*, **19**, 669-675.

A New Cytotoxic Cyclolanostane Triterpenoid Xyloside from <i>Souliea vaginata</i> Haifeng Wu, Zhixin Yang, Qiru Wang, Nailiang Zhu, Xudong Xu, Qiongyu Zou and Yulian Tang	229
Polyhydroxy Sterols Isolated from the Red Sea Soft Coral <i>Lobophytum crassum</i> and their Cytotoxic Activity Elsayed A. Aboutabl, Nabil M. Selim, Shadia M Azzam, Camilia G. Michel, Mohamed F. Hegazy, Abdelhamid M. Ali and Ahmed A. Hussein	233
<i>Raphanus sativus</i> Sprout Causes Selective Cytotoxic Effect on p53-Deficient Human Lung Cancer Cells <i>in vitro</i> Jiwon Baek, Hyun-Soo Roh, Chang-Ik Choi, Kwan-Hyuck Baek and Ki Hyun Kim	237
Preparative and Rapid Purification of Saponins from <i>Asparagus racemosus</i> Root by High Performance Centrifugal Partition Chromatography Churanya Onlom, Yi Yang, Haji A. Aisa, Neti Woranuch, Watoo Phrompittayarat, Waraporn Putalun and Kornkanok Inghaninan	241
AKR1C3 Inhibitory Potency of Naturally-occurring Amaryllidaceae Alkaloids of Different Structural Types Daniela Hulcová, Kateřina Breiterová, Lucie Zemanová, Tomáš Siatka, Marcela Šafratová, Nina Vaněčková, Anna Hošťálková, Vladimír Wsól and Lucie Cahlíková	245
Product Selectivity of Esterification of L-Aspartic Acid and L-Glutamic Acid Using Chlorotrimethylsilane Tomohiro Takaishi, Minoru Izumi, Ryo Ota, Chieri Inoue, Hiromasa Kiyota and Koichi Fukase	247
New Cyclopentyl Fatty Acid and Cyanohydrin Glycosides from Fruits of <i>Hydnocarpus hainanensis</i> Thanh Tra Nguyen, Bich Ngan Truong, Huong Doan Thi Mai, Marc Litaudon, Van Hung Nguyen, Thao Do Thi, Van Hieu Tran, Dang Thach Tran, Van Minh Chau and Van Cuong Pham	251
Chemical Constituents of the Roots and Rhizomes of <i>Saposhnikovia divaricata</i> and their Cytotoxic Activity Akihito Yokosuka, Satoru Tatsuno, Takuma Komine and Yoshihiro Mimaki	255
Cytotoxic Activity of Compounds from <i>Styrax obassia</i> Thao Quyen Cao, Bo Mi Lee, Yeon Woo Jung, Van Thu Nguyen, Jeong Ah Kim and Byung Sun Min	259
Isolation and Characterization of Antiangiogenesis Compounds from the Fungus <i>Aspergillus terreus</i> Associated with <i>Apostichopus japonicus</i> Using Zebrafish Assay Jun Qi, Bo Zhao, Peipei Zhao, Airong Jia, Yonggang Zhang, Xin Liu, Changheng Liu, Lixin Zhang and Xuekui Xia	261
Adsorptive Property of Food Materials and Chemicals to Cesium and Strontium Yuya Uehara, Kana Ishizuka, Yuko Shimamura, Michiko Yasuda, Kayoko Shimoi and Shuichi Masuda	263
Germination and Clonal Propagation of the Endemic Shrub <i>Corema album</i>, a Vulnerable Species with Conservation Needs and Commercial Interest Leonor Álvarez-Cansino, María Zunzunegui and Mari Cruz Díaz Barradas	267
Essential Oil Profile, Phenolic Content and Antioxidant Activity of <i>Geranium kikianum</i> Sanja Čavar Zeljković, Kit Tan, Sonja Siljak-Yakovlev and Milka Maksimović	273
Integrated Analysis of the Bark Oil from <i>Cinnamosma madagascariensis</i> by GC(RI), GC-MS and NMR. ¹³C NMR data of Cyclocopacamphene and Cyclosativene Gabriel Garcia, Delphin Rabehaja, Julie-Marie Charmillon, Panja Ramanoelina, Joseph Casanova and Félix Tomi	277
Synergistic Activity of Essential Oils from Herbs and Spices Used on Meat Products against Food Borne Pathogens Juan García-Díez, Joana Alheiro, Ana Luisa Pinto, Virgilio Falco, Maria João Fraqueza and Luís Patarata	281
Composition, <i>in vitro</i> Cytotoxicity, Anti-mildew and Anti-wood-decay Fungal Activities of the Fruit Essential Oil of <i>Liquidambar formosana</i> from Taiwan Yu-Chang Su and Chen-Lung Ho	287
Chemical Composition and Antibacterial Activity of the Bulgarian Endemic Species <i>Achillea thracica</i> from its Natural Habitat, and <i>in vitro</i> Propagated and <i>ex vitro</i> Established Plants Gordana S. Stojanović, Veneta Kapchina-Toteva, Mariq Angelova Rogova, Snežana Č. Jovanović, Zhenya Yordanova and Bojan K. Zlatković	291
<u>Accounts/Reviews</u>	
<i>In vitro</i> and <i>in vivo</i> Methods for the Evaluation of Natural Products against Dermatophytes Lily A. Velázquez-Dávila, Sergio A. Galindo-Rodríguez, Luis A. Pérez-López, M. Gloria González-González and Rocío Álvarez-Román	293
Chemistry and Pharmacology of <i>Tinospora cordifolia</i> Deepika Singh and Prabir K Chaudhuri	299
Perspectives of the <i>Apiaceae</i> Hepatoprotective Effects – A Review Milica G. Acimović and Nataša B. Milić	309

Natural Product Communications 2017

Volume 12, Number 2

Contents

NPC-CMAPSEEC Issue

Original Paper

- Phytochemical Profile of *Inula britannica* from Bulgaria**
Victoria Ivanova, Antoaneta Trendafilova, Milka Todorova, Kalina Danova and Dimitar Dimitrov 153
- Production of Δ^7 -Sterols from *In Vitro* Root Cultures of Endangered *Gypsophila trichotoma***
Petranka Zdraveva, Pavlinka Popova, Aleksandar Shkondrov, Iliana Krasteva and Iliana Ionkova 155
- Evaluation of Glaucine Content in Bulgarian Black Sea Coast Localities of *Glaucium flavum* Cranz. (Papaveraceae)**
Iva Doycheva, Stefan Philipov and Marina Stanilova 157
- Crataegus orientalis* Leaves and Berries: Phenolic Profiles, Antioxidant and Anti-inflammatory Activity**
Katarina P. Šavikin, Dijana B. Krstić-Milošević, Nebojša R. Menković, Ivana N. Beara, Zorica O. Mrkonjić and Dejan S. Pljevljakušić 159
- Analysis of Antioxidant Polyphenols in Loquat Leaves using HPLC-based Activity Profiling**
Izabela Nawrot-Hadzik, Sebastian Granica, Renata Abel, Hanna Czapor-Irzabek and Adam Matkowski 163
- LC/DAD/MSⁿ and ICP-AES Assay and Correlations between Phenolic Compounds and Toxic Metals in Endemic *Thymus alsarensis* from the Thallium Enriched Allchar Locality**
Jasmina Petreska Stanoeva, Marina Stefova, Katerina Bačeva Andonovska and Trajče Stafilov 167
- Ultrasound and Microwave-Assisted Extraction of *Inula helenium* Roots**
Nadezhda Petkova, Ivan Ivanov, Radka Vrancheva, Panteley Denev and Atanas Pavlov 171
- ADME/Tox Properties and Biochemical Interactions of Silybin Congeners: *In silico* Study**
Antonia Diukendjjeva, Merilin Al Sharif, Petko Alov, Tania Pencheva, Ivanka Tsakovska and Ilza Pajeva 175
- Comparative Study of Naphthoquinone Contents of Selected Greek Endemic Boraginaceae Plants - Antimicrobial Activities**
Teisa Tufa, Harilaos Damianakos, Konstantia Graikou and Ioanna Chinou 179
- Effects of Gamma-Irradiation on the Antioxidant Potential of Traditional Bulgarian Teas**
Michał Adam Janiak, Adriana Slavova-Kazakova, Magdalena Karamać, Vessela Kancheva, Anastasiya Terzieva, Milena Ivanova, Tsvetelin Tsrunchev and Ryszard Amarowicz 181
- Microelements and Heavy Metals Content in Frequently Utilized Medicinal Plants Collected from the Power Plant Area**
Aleksandra Stanojković-Sebić, Jelena Maksimović, Zoran Dinić, Dobrovojo Poštić, Renata Iličić, Aleksandar Stanojković and Radmila Pivić 185
- Molecular Characterization of *Verbascum anisophyllum* (Scrophulariaceae) Genetic Resources Through Inter-Simple Sequence Repeat (ISSR) Markers**
Galya Petrova, Stefan Petrov and Svetlana Bancheva 189
- Effect of Drought and Salinity on Volatile Organic Compounds and Other Secondary Metabolites of *Citrus aurantium* Leaves**
Sarrou Eirini, Chatzopoulou Paschalina, Therios Ioannis and Dimassi-Theriou Kortessa 193
- Insights into the Essential Oil Compositions of Brazilian Red and Taiwanese Green Propolis**
Boryana Trusheva, Daniela Ivanova, Milena Popova and Vassya Bankova 197
- Essential Oil Content, Composition and Bioactivity of Juniper Species in Wyoming, United States**
Valtcho D. Zheljatzkov, Tess Astatkie, Ekaterina A. Jeliatzkova, Bonnie Heidel and Lyn Ciampa 201
- Chemical Composition and Antibacterial Activity of *Angelica archangelica* Root Essential Oil**
Milica G. Aćimović, Snežana Đ. Pavlović, Ana O. Varga, Vladimir M. Filipović, Mirjana T. Cvetković, Jovana M. Stanković and Ivana S. Čabarkapa 205

Original Paper

- Gas Chromatography-Mass Spectrometry (GC-MS) Combined with Retention Index Prediction for the Rapid Identification of Halogenated monoterpenes from a Namibian *Plocamium* species**
Stefan Louw, Lineekela Kandjengo and Michael G. Knott 207
- Rapid and Efficient Extraction and HPLC Analysis of Sesquiterpene Lactones from *Aucklandia lappa* Root**
Clizia Guccione, Giorgia Ros, Sandra Gallori, Maria Camilla Bergonzi and Anna Rita Bilia 213
- Antimalarial Activity of some Kaurenes**
Thayded Villasmil, Julio Rojas, Rosa Aparicio, Neira Gamboa, Maria Eugenia Acosta, Juan Rodrigues and Alfredo Usubillaga 217
- Briarenol B, a New Polyoxygenated Briarane from the Octocoral *Briareum excavatum***
Mu-Jang Li, Yin-Di Su, Zuo-Jian Liao, Zhi-Hong Wen, Jui-Hsin Su, Yang-Chang Wu and Ping-Jyun Sung 221
- Triterpenes with Anti-invasive Activity from Sclerotia of *Inonotus obliquus***
Kaori Ryu, Seikou Nakamura, Souichi Nakashima, Masaaki Aihara, Masashi Fukaya, Junko Iwami, Yasunobu Asao, Masayuki Yoshikawa and Hisashi Matsuda 225

Continued inside backcover